

Healthy Infants and Young Children Milk

Applicable to: Nurses, Physicians and Other Health Professionals

Recommendations

- Human milk is the optimal milk for infants and is recommended for up to two years and beyond.
- Pasteurized 3.25% (homogenized) milk may be introduced to healthy term infants at 9–12 months of age and continued throughout the second year of life.
- Whole, 3.25% or greater, pasteurized goat's milk fortified with vitamin D may be used as an alternative to cow's milk after 9–12 months of age.
- Children one year of age and older should be offered 2 cups (500 mL) of milk each day.
- Children two years of age and older can be offered lower fat milk (skim, 1% or 2%).
- Soy, rice, almond, and other plant-based beverages are not suitable alternatives to milk for children under two years of age.
- For children two years of age and older not consuming milk, Canada's food guide recommends a fortified soy beverage. If parents or caregivers choose another plant-based beverage as a milk alternative for their child, they should look for a product that:
 - provides at least 6 g of protein per 1 cup (250 mL);
 - provides at least 30% daily value of calcium and vitamin D per 1 cup (250 mL); and
 - contains less than 10 g of sugar per 1 cup (250 mL).

Notice, March 2022: There has been a practice change regarding the recommendations for consumption of plant-based beverages for infants and young children. Based on new labelling guidelines, the % daily values for calcium are now **23% (300 mg)** and **10% (80 IU)** for vitamin D per 1 cup (250 mL). In addition, parents or caregivers should look for a product that is fortified or enriched and contains less than **15% daily value** for sugar (<15 g) per 1 cup (250 mL). Please see the [Plant-based Beverages](#) for Infants and Young Children Nutrition Guideline for more information.

- A child under two years of age who is not drinking human milk, 3.25% milk, or commercial infant formula may benefit from a referral to a dietitian.
- Toddler formulas which are marketed as an alternative or complement to human milk or fluid milk are not necessary for young children.



Nutrition Guideline

Healthy Infants and Young Children

Milk

Health Benefits

This guideline will provide information on and considerations for the use of fluid milks for infants and young children. This information will assist health professionals when educating parents and caregivers on appropriate milk choices for infants and young children to help support healthy growth and development. Fluid milks contain nutrients such as protein, calcium, vitamin D, phosphorus, potassium, magnesium, and vitamins A, B₁₂, and riboflavin which support growth and development including building strong bones and teeth.¹

Definitions

Young infant: Birth to six months

Older infant: 6–12 months

Young child: 12–24 months

Infant formula: Food intended for young infants until the introduction of appropriate complementary feeding.

Follow-up formula: Formula marketed for infants six months and older when appropriate complementary feeding is introduced.

Milk: The fluid secreted by female mammals to feed their young. For this guideline, human milk and fortified cow or goat milk are referred to as “milk”.

Plant-based beverage: A beverage made from legumes, nuts, cereal grains, or potatoes.² Common examples are soy, almond, and rice beverages.

Toddler formula (Growing up milk, milk beverage, milk drink): A supplement marketed as an alternative or complement to human milk and cow’s milk for young children.

Nutrition Guideline

Healthy Infants and Young Children

Milk

Key Questions

At what age can cow's milk be introduced?

At 9–12 months of age, healthy term infants can be offered pasteurized, 3.25% (homogenized) milk.³ This can also be in the form of reconstituted powder 3.25% milk or evaporated canned 3.25% milk.² Preterm infants should not receive 3.25% milk until they are 12 months corrected age.^{4–6} To reduce the risk of iron deficiency, older infants need to consume iron-rich foods at most meals before 3.25% milk is offered.⁷ It is best to offer milk at meals and snacks in an open cup. Sipping on milk between meals may cause early childhood caries.^{8,9}

Cow's milk to drink is not appropriate for infants younger than nine months of age as it:

- is low in iron^{3,10}
- may displace iron-rich foods³
- can inhibit iron absorption³
- is low in essential fatty acids¹⁰
- contains a less digestible form of protein¹⁰
- is associated with occult blood loss in stool in infants under six months of age^{11–14}

After six months of age, milk can be used in baking or cooking, and milk products such as cheese and yogurt can be introduced along with a variety of iron-rich foods.³

For infants younger than nine months of age not receiving human milk, commercial infant formula will meet the nutritional needs of infants and is the only recommended alternative to human milk.³

For more information on infant formula or vitamin D, refer to the [Nutrition Guidelines: Infant Formulas or Vitamin D](#).

If an older infant or child is breastfed, can you offer additional milk to drink?

Breastfeeding can be continued for up to two years of age and beyond, as long as the mother and child want to continue.¹⁵ Older breastfed infants and children not consuming foods or drinks from the Milk and Alternatives food group (e.g. yogurt and cheese) may not be getting enough calcium and vitamin D. Nutrient adequacy will be dependent on the frequency and volume of human milk consumed. Individual guidance from a dietitian may be warranted.

If a parent or caregiver chooses to offer 3.25% milk to a breastfed child, then it is best to offer it at meals and snacks and in an open cup. For children one year and older, the total amount from both milk and human milk should be no more than 2 cups (500 mL) to prevent iron deficiency anemia.⁷ Children who are overly reliant on milk or human milk may consume insufficient amounts of complementary foods which can increase the risk of nutrient deficiencies, such as iron.^{16,17}

Health professionals can support human milk feeding of the older infant and young child by continuing to promote this practice as the normal way of feeding.³

Nutrition Guideline

Healthy Infants and Young Children

Milk

What are the considerations around the consumption of goat's milk?

Pasteurized, whole goat's milk (3.25% or greater) fortified with vitamin D can be offered as an alternative to cow's milk once an infant is 9–12 months of age and eating iron-rich foods at most meals.³ Similar to cow's milk, lower-fat goat's milk are not appropriate before two years of age. Goat's milk poses the same risks for the development of iron deficiency as cow's milk when consumed in excessive amounts.^{3,18} Just like cow's milk, goat's milk is not appropriate for infants younger than nine months as it:

- is low in iron
- is low in essential fatty acids and other essential nutrients
- contain a less-digestible form of protein³

Goat's milk is not a recommended alternative to cow's milk for cow's milk allergy or for lactose intolerance. Cow's and goat's milk are very similar in composition in terms of their protein, fat, and carbohydrate content.¹⁹ Due to the similarity in proteins, older infants and young children who have a food allergy to cow's milk protein are also likely to have an allergic reaction to goat's milk.³

Although both goat's milk and cow's milk contain comparable amounts of lactose, goat's milk contains slightly lower levels.¹⁹ Goat's milk contains smaller fat droplets which may ease digestibility, but this has not been proven.²⁰

Goat's milk is significantly lower in folic acid unless it is fortified. However, at 9–12 months of age, older infants will receive folate through complementary foods such as vegetables (especially dark green vegetables), fruits, beans, poultry and meat, eggs, seafood, and grains.²¹

Health Canada recommends choosing a goat's milk fortified with vitamin D, however, fortification with vitamin D is not mandatory in Canada.³ Currently, there are no goat's milk products fortified with vitamin D available for purchase in Alberta, therefore, parents or caregivers who offer non-fortified whole goat's milk should be advised that their child may not meet their daily vitamin D requirements. Parents or caregivers may wish to ask their local store to use a supplier that offers fortified goat's milk. Parents or caregivers offering non-fortified goat's milk should consult with their child's doctor or an RD to see if additional vitamin D supplementation is required. The nutrient content for both milks are otherwise similar. Refer to Table 1.

Nutrition Guideline

Healthy Infants and Young Children

Milk

Table 1. Comparison of Cow's and Goat's Milk Nutrient Content per 1 cup (250 mL)¹⁹

Nutrient	Cow's milk	Goat's milk	
	3.25% (Homogenized)	Fortified (Whole)	Unfortified (Whole)
Energy (calories)	157	178	178
Protein (g)	8	9	9
Carbohydrate (g)	12	11	11
Fat (g)	8	11	11
Calcium (mg)	291	345	345
Iron (mg)	0.1	0.1	0.1
Vitamin A (RAE-mcg)	119	147	147
Vitamin D (IU)	103	100	0.3
Folic Acid (DFE – ug)	13	37	2

RAE: Retinol Activity Equivalents

DFE: Dietary Folate Equivalents

How much milk should children drink?

It is recommended that children one year of age and older drink 2 cups (500 mL) of milk each day.³ Cow's milk provides a child with calcium, vitamin D, protein, and other important vitamins and minerals. The recommendation of 2 cups (500 mL) per day provides approximately 200 IU of vitamin D and 600 mg of calcium per day.¹ Consumption of more than 2 cups (500 mL) a day has been identified as a risk factor associated with iron deficiency anemia^{22–24} as children may fill up on milk, displacing iron-rich foods in their diet. Offering milk in an open cup may help to avoid excess consumption.^{3,25}

For children who are breastfed at one to two years of age, approximately $\frac{1}{3}$ of a child's energy should generally come from human milk and the remaining $\frac{2}{3}$ from complementary foods.¹⁶ Children who are overly reliant on human milk as their main source of nutrition may consume insufficient amounts of complementary foods,¹⁶ which can put young children at risk of nutrient deficiencies, such as iron. For children consuming a combination of human milk, cow's milk and/or formula, the total amount of milk from all sources should be no more than 2 cups (500 mL) per day. Cow or goat's milk and formula should be offered in an open cup to avoid excess consumption.²⁵

For more information on iron and excessive milk consumption, refer to the Nutrition Guidelines: [Iron and Healthy Feeding Relationship](#).

For information related to calcium and vitamin D, refer to the Nutrition Guideline: [Calcium and Vitamin D](#)

Nutrition Guideline

Healthy Infants and Young Children

Milk

At what age can a child drink lower fat milk?

Children two years of age and older can be offered lower fat milk choices (skim, 1% or 2% or fortified soy beverage) to drink. This transition to lower fat choices can be gradual.²⁶

Skim milk is an inappropriate choice for children younger than two years.³ Lower fat milks are low in energy and may cause growth faltering.³ If a child is offered 1% or 2% milk before two years of age, a referral to a dietitian or a physician/pediatrician is recommended to ensure that the child is growing well.³ In addition, advise parents and caregivers to provide higher-fat, nutritious foods that are an important source of energy such as human milk, 3.25% milk, cheese, avocado, nut butters, and some fish such as salmon or trout.²⁷

What if a child does not drink milk?

Older infants and young children who do not receive human milk and who refuse to drink fluid milks can still receive milk as part of other foods (e.g. in soups, smoothies, added to cereal). Encourage parents and caregivers to continue to offer small quantities of milk at meals and snacks²⁸ as a child's vitamin D needs may not be met with milk alternatives such as yogurt, cheese, and drinkable yogurt.²⁸ These products may not be fortified with sufficient vitamin D.

For children 12–24 months who are unable to drink milk due to an allergy or intolerance, or follow a vegetarian or vegan lifestyle, advise parents and caregivers to offer commercial infant follow-up soy formula.³ Over two years of age, if milk is not consumed, Canada's food guide recommends a fortified soy beverage.²

When milk products are removed from the diet, protein, calcium, and vitamin D intakes are decreased and must be replaced through other sources. A Registered Dietitian (RD) can assess if a child is meeting their nutrient requirements and can suggest ways to adjust dietary intake.

Notice, March 2022: There has been a practice change regarding the recommendations for consumption of soy follow-up formula. If infants are drinking a soy formula, continue to offer this formula until 24-months of age along with calcium rich complementary foods. In previous years, follow-up soy formula was recommended between 12–24 months of age; however, they are no longer available. Please see the [Plant-based Beverages](#) for Infants and Young Children Nutrition Guideline for more information.

For more information, refer to the Nutrition Guidelines: [Infant Formulas and Plant-Based Beverages](#)

Nutrition Guideline

Healthy Infants and Young Children

Milk

Are plant-based beverages appropriate alternatives for milk in children?

Children Under Two Years of Age

Plant-based beverages are not appropriate in the first two years of life. These beverages do not support proper growth and development as they are lower in protein, fat, and calories, and may not be fortified with calcium and vitamin D.^{3,29} Infants who are drinking a commercial soy formula should drink a **follow-up** soy formula from 12–24 months of age.

If fortified soy beverage is being offered occasionally as a complementary food (i.e. in addition to human milk or cow's milk as the main milk source) unflavoured, fortified soy beverage is recommended.³⁰

Notice, March 2022: There has been a practice change regarding the recommendations for consumption of soy follow-up formula. If infants are drinking a soy formula, continue to offer this formula until 24 months of age along with calcium rich complementary foods. In previous years, follow-up soy formula was recommended between 12–24 months of age; however, they are no longer available. Please see the [Plant-based Beverages](#) for Infants and Young Children Nutrition Guideline for more information.

Children Two Years of Age and Older

If milk is not consumed, Canada's food guide recommends a fortified soy beverage.² If parents or caregivers choose a plant-based beverage as a milk alternative for their child age two years and older, they should look for a product that:

- is labelled as fortified or enriched;³¹
- provides at least 6 g of protein per 1 cup (250 mL),³¹
- provides at least 30% daily value of calcium and vitamin D per 1 cup (250 mL); and
- contains less than 10 g of sugar per 1 cup (250 mL).³²

Notice, March 2022: There has been a practice change regarding the recommendations for consumption of plant-based beverages for infants and young children. Based on new labelling guidelines the % Daily Values for Calcium are now **23% (300 mg)** and **10% (80 IU)** for vitamin D per 1 cup (250 mL). In addition, parents or caregivers should look for a product that is fortified or enriched and contains less than **15% Daily Value** for sugar (<15 g) per 1 cup (250 mL). Please see the [Plant-based Beverages](#) for Infants and Young Children Nutrition Guideline for

For more information on plant-based beverages, refer to the Nutrition Guideline: [Plant-based Beverages](#).

For more information on soy formulas, refer to the [Infant Formulas for Healthy Term Infants Compendium and Summary Sheet](#).

Is unpasteurized milk safe?

Raw or unpasteurized milk is not recommended due to the risk of foodborne illness. The major symptoms resulting from drinking unpasteurized milk include:

- Vomiting
- Diarrhea (which may be bloody)
- Stomach cramps or abdominal pain
- Life-threatening kidney failure³³

Nutrition Guideline

Healthy Infants and Young Children

Milk

Pasteurization is a process that kills harmful bacteria that can cause foodborne illness while keeping the nutritional properties of milk intact.³⁴ Therefore, Food and Drug Regulations require that all milk available for sale in Canada be pasteurized.³⁴

Although it is illegal to sell unpasteurized milk in Canada, some people access raw milk from farms. It is important to note that although some Canadians prefer raw milk because they believe it is healthier, any possible benefits are far outweighed by the serious risk of illness from drinking raw milk.³⁴

Although it is not recommended, if a parent or caregiver chooses to provide raw milk to their child, it should be pasteurized before it is consumed. Parents and caregivers should be advised that it will also not contain the vitamin D that is added to commercial Canadian cow's milk. Information on home pasteurization can be found at: <https://myhealth.alberta.ca/Alberta/Pages/how-to-pasteurize-milk.aspx>

Does milk contain hormones and antibiotics?

Milk produced and sold in Canada does not contain added hormones. The growth hormone recombinant bovine somatotropin (rBST) which increases milk production is not approved for use in Canada. Milk produced in the United States of America may contain rBST.³⁵

Milk produced and sold in Canada does not contain antibiotics. Antibiotics are used to treat diseases in animals. If a cow is sick and needs medicine such as antibiotics, the milk she makes while she's on medications and for a period of time after must be discarded.³⁶

What is the difference between organic and conventional milk?

The main difference between organic and conventional milk production are the farming practices. Organic producers do not use synthetic pesticides, herbicides, fungicides or nitrogen-based fertilizers.³⁷ Canadian organic production follows strict standards as set out in the Canadian Organic Standards.³⁸ Organically raised dairy cows are not confined to pens and are given living conditions that allow them access to a pasture during the grazing season, as well as access to open air at other times weather permitting.³⁸ Similar to conventionally raised cows, the milk from organic cows that are being treated for illness must be discarded while they are sick.^{36,38} Organic cows that are treated for illness are typically kept out of production longer after treatment and the withdrawal period is twice the regular period or at least two weeks, whichever is longer. Treatment with antibiotics results in a milk withdrawal period of at least 30 days, after the last day of a course of treatment, or a withholding period that is twice the label requirement, whichever is longer. If an animal is treated with antibiotics more than once in a year, it is permanently removed from production.³⁹ Evidence suggests organically produced food may contain less exposure to pesticides but does not suggest they are more nutritious compared to conventional foods.⁴⁰

What are the considerations for the use of toddler formulas?

Toddler formulas are also referred to as toddler nutritional supplements or more commonly as 'Growing-up milk' or 'toddler milk' such as Enfagrow A+™, Similac® Go & Grow®, Good Start® 3 Toddler Transition, Gerber® Graduates Toddler Drink and store brands such as President's Choice® Toddler and Parent's Choice™ Toddler. They are marketed as an alternative or

Nutrition Guideline

Healthy Infants and Young Children

Milk

complement to human milk or fluid milk for children older than 12 months of age.^{41–43} Toddler milks are not necessary or recommended for healthy toddlers. Consuming toddler milks can displace human milk and other healthy foods in the diet. They can be more costly, have added sugars and do not provide additional benefits when compared to human milk or standard, age-appropriate milk. Some potential risks of using toddler milks include over and under dilution due to improper mixing.⁴⁴

It is recommended that young children (toddlers) eat a variety of healthy foods to supply the majority of their nutrients, rather than relying on fortified toddler formulas.⁴¹ Nutrient needs can generally be met by following Canada's food guide.²

If parents and caregivers have further questions on the use of toddler formulas, advise that they speak with an RD for further assessment.

Are flavoured milks appropriate to drink?

Flavoured milks such as chocolate milk contain the same nutrients as plain cow's milk but are usually lower fat and contain added sugars. Plain, 3.25% milk is recommended over flavoured milks, such as chocolate milk, or flavoured soy-based beverages, such as chocolate and vanilla, which contain added sugar.¹⁰

Are there any resources related to feeding healthy infants and children that I can use with my patients?

For nutrition resources visit Nutrition Education Materials at www.albertahealthservices.ca/nutrition/Page11115.aspx and click on **Infants or Children/Adolescents**.

For more information related to healthy infants and children see [Healthy Parents Healthy Children](#).

Nutrition Guideline

Healthy Infants and Young Children

Milk

References

1. Institute of Medicine. Dietary reference intakes for calcium and vitamin D. Washington, DC: National Academies Press; 2011.
2. Health Canada. Canada's Food Guide [Internet]. 2021 [cited 2021 May 3]. Available from: <https://food-guide.canada.ca/en/>
3. Health Canada. Nutrition for Healthy Term Infants: Recommendations from Six to 24 Months [Internet]. Vol. 2019. 2014. Available from: <https://www.canada.ca/en/health-canada/services/canada-food-guide/resources/infant-feeding/nutrition-healthy-term-infants-recommendations-birth-six-months/6-24-months.html>
4. American Academy of Pediatrics. Age Terminology During the Perinatal Period. *Pediatrics* [Internet]. 2004 [cited 2022 May 2];114(5):1362–4. Available from: <https://publications.aap.org/pediatrics/article/114/5/1362/67715/Age-Terminology-During-the-Perinatal-Period>
5. Canadian Pediatric Society Nutrition Committee. Nutrient needs and feeding of premature infants. Nutrition Committee, Canadian Paediatric Society - PubMed. *Can Med Assoc J* [Internet]. 1995 [cited 2022 May 2];152(11):1765–85. Available from: <https://pubmed.ncbi.nlm.nih.gov/7773894/>
6. Rao R, Georgieff MK. Iron therapy for preterm infants. *Clin Perinatol* [Internet]. 2009 Mar [cited 2022 May 2];36(1):27–42. Available from: <https://pubmed.ncbi.nlm.nih.gov/19161863/>
7. Agostoni C, Turck D. Is cow's milk harmful to a child's health? *J Pediatr Gastroenterol Nutr* [Internet]. 2011 Dec [cited 2022 May 2];53(6):594–600. Available from: <https://pubmed.ncbi.nlm.nih.gov/21921812/>
8. American Dental Association. From baby bottle to cup: Choose training cups carefully, use them temporarily. *J Am Dent Assoc* [Internet]. 2004 Mar 1 [cited 2022 May 2];135(3):387. Available from: <http://jada.ada.org/article/S0002817714612113/fulltext>
9. Çolak H, Dülgergil Ç, Dalli M, Hamidi M. Early childhood caries update: A review of causes, diagnoses, and treatments. *J Nat Sci Biol Med* [Internet]. 2013 Jan [cited 2022 May 2];4(1):29. Available from: </pmc/articles/PMC3633299/>
10. Health Canada, Canadian Pediatric Society Nutrition Committee, Dietitians of Canada, Breastfeeding Committee for Canada. Nutrition for healthy term infants: recommendations from birth to six months [Internet]. Vol. 2017. 2015. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/healthy-eating/infant-feeding/nutrition-healthy-term-infants-recommendations-birth-six-months.html>
11. Bondi S, Lieuw K. Excessive cow's milk consumption and iron deficiency in toddlers: two unusual presentations and review. *Infant, Child Adolesc Nutr*. 2009;1(3):133–9.
12. Fernandes SMR, De Morais MB, Amancio OMS. Intestinal blood loss as an aggravating factor of iron deficiency in infants aged 9 to 12 months fed whole cow's milk. *J Clin Gastroenterol* [Internet]. 2008 Feb [cited 2022 May 2];42(2):152–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/18209584/>
13. Fomon SJ, Nelson SE, Serfass RE, Ziegler EE. Absorption and Loss of Iron in Toddlers Are Highly Correlated. *J Nutr* [Internet]. 2005 Apr 1 [cited 2022 May 2];135(4):771–7. Available from: <https://academic.oup.com/jn/article/135/4/771/4663751>
14. Michaelsen KF. Cows' Milk in Complementary Feeding. *Pediatrics* [Internet]. 2000 Nov 1 [cited 2022 May 2];106(Supplement_4):1302–3. Available from: /pediatrics/article/106/Supplement_4/1302/65869/Cows-Milk-in-Complementary-Feeding
15. World Health Organization. Infant and young child feeding [Internet]. [cited 2022 May 2]. Available from: <https://www.who.int/news-room/fact-sheets/detail/infant-and-young-child-feeding>
16. Dewey KG, Brown KH. Update on technical issues concerning complementary feeding of young children in developing countries and implications for intervention programs. *Food Nutr Bull* [Internet]. 2003;24(1):5–28. Available from: <http://ahs.idm.oclc.org/login?url=http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med4&AN=12664525>
<http://sfxhosted.exlibrisgroup.com/hinc?sid=OVID:medline&id=pmid:12664525&id=doi:&issn=0379-5721&isbn=&volume=24&issue=1&spage=5&pages=>
17. Dube K, Schwartz J, Mueller MJ, Kalhoff H, Kersting M. Iron intake and iron status in breastfed infants during the

Nutrition Guideline

Healthy Infants and Young Children

Milk

- first year of life. Clin Nutr [Internet]. 2010 Dec [cited 2022 May 2];29(6):773–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/20627488/>
18. Centers for Disease Control and Prevention (CDC). Recommendations to Prevent and Control Iron Deficiency in the United States [Internet]. [cited 2022 May 2]. Available from: <https://www.cdc.gov/mmwr/preview/mmwrhtml/00051880.htm>
 19. Health Canada. Canadian Nutrient File. Version 2015 [database on the internet] [Internet]. Vol. 2018. 2016. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/healthy-eating/nutrient-data.html#wb-cont>
 20. Raynal-Ljutovac K, Lagriffoul G, Paccard P, Guillet I, Chilliard Y. Composition of goat and sheep milk products: An update. Small Rumin Res [Internet]. 2008 Sep [cited 2022 May 2];1(79):57–72. Available from: <https://www.infona.pl/resource/bwmeta1.element.elsevier-e2d309d2-7e8c-36cb-8b99-bbb7f4af93ec>
 21. Dietitians of Canada. Food sources of folate. Unlock Food [Internet]. [cited 2022 May 2]. Available from: <https://www.unlockfood.ca/en/Articles/Vitamins-and-Minerals>
 22. Domellöf M, Braegger C, Campoy C, Colomb V, Decsi T, Fewtrell M, et al. Iron requirements of infants and toddlers. J Pediatr Gastroenterol Nutr [Internet]. 2014 Jan [cited 2022 May 2];58(1):119–29. Available from: <https://pubmed.ncbi.nlm.nih.gov/24135983/>
 23. Parkin PC, Degroot J, Maguire JL, Birken CS, Zlotkin S. Severe iron-deficiency anaemia and feeding practices in young children. Public Health Nutr [Internet]. 2016 Mar 1 [cited 2022 May 2];19(4):716–22. Available from: <https://pubmed.ncbi.nlm.nih.gov/26027426/>
 24. Paoletti G, Bogen DL, Ritchey AK. Severe iron-deficiency anemia still an issue in toddlers. Clin Pediatr (Phila) [Internet]. 2014;53(14):1352–8. Available from: <http://www.sagepub.com/journalsProdDesc.nav?prodId=Journal201788>
<http://ahs.idm.oclc.org/login?url=http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed12&AN=2014922357> <http://sfxhosted.exlibrisgroup.com/hinc?sid=OVID:embase&id=pmid:&i>
 25. Maguire JL, Lebovic G, Kandasamy S, Khovratovich M, Mamdani M, Birken CS, et al. The relationship between cow's milk and stores of vitamin D and iron in early childhood. Pediatrics. 2013 Jan;131(1):144.
 26. American Dietetic Association. Position of the American Dietetic Association and Dietitians of Canada: dietary fatty acids. J Am Diet Assoc. 2007;107:1599–611.
 27. Health Canada. Infant nutrition [Internet]. [cited 2022 May 2]. Available from: <https://www.canada.ca/en/health-canada/services/infant-care/infant-nutrition.html>
 28. Satter E. Child of Mine: Feeding With Love and Good Sense [Internet]. [cited 2022 May 2]. Available from: <https://www.ellynsatterinstitute.org/product/child-of-mine/>
 29. Health Canada. Healthy eating resources - Canada's Food Guide [Internet]. [cited 2022 May 2]. Available from: <https://food-guide.canada.ca/en/healthy-eating-resources/>
 30. Dunham L, Kollar LM. Vegetarian eating for children and adolescents. J Pediatr Health Care [Internet]. 2006 Jan [cited 2022 May 2];20(1):27–34. Available from: <https://pubmed.ncbi.nlm.nih.gov/16399477/>
 31. Health Canada. Government notices – Department of Health, Food and Drugs Act. Interim marketing authorization, Appendix F – interim marketing authorization for amendments to plant-based beverages. [Internet] [Internet]. Vol. 2018. 1997. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/legislation-guidelines/policies/interim-policy-on-use-expired-interim-marketing-authorizations-related-food-fortification.html#a61>
 32. Government of Alberta, Alberta Health. Alberta Nutrition Guidelines for Children and Youth A Childcare, School and Recreation/Community Centre Resource Manual. 2012;
 33. MyHealthAlberta. How to pasteurize milk at home [Internet]. [cited 2022 May 2]. Available from: <https://myhealth.alberta.ca/Alberta/Pages/how-to-pasteurize-milk.aspx>
 34. Health Canada. Risks of Drinking Raw Milk [Internet]. [cited 2022 May 2]. Available from: <https://recalls-rappels.canada.ca/en/alert-recall/risks-drinking-raw-milk-1>
 35. Health Canada. Questions and Answers - Hormonal Growth Promoters [Internet]. [cited 2022 May 2]. Available from: <https://www.canada.ca/en/health-canada/services/drugs-health-products/veterinary-drugs/factsheets-faq/hormonal-growth-promoters.html>

Nutrition Guideline

Healthy Infants and Young Children

Milk

36. Dietitians of Canada. Food Safety - Hormone and Antibiotic Background. In: Practice-based Evidence in Nutrition [PEN] [Internet]. 2011 [cited 2022 May 2]. Available with subscription.
37. Government of Canada. Organic Dairy Production [Internet]. [cited 2022 May 2]. Available from: <http://www.omafra.gov.on.ca/english/livestock/dairy/facts/10-087.htm>
38. Government of Canada. Organic dairy industry in Canada [Internet]. [cited 2022 May 2]. Available from: <https://publications.gc.ca/site/eng/393195/publication.html>
39. Government of Canada. Organic production systems : general principles and management standards. [Internet]. [cited 2022 May 2]. Available from: <https://publications.gc.ca/site/eng/9.854643/publication.html>
40. Smith-Spangler C, Brandeau ML, Hunter GE, Clay Bavinger J, Pearson M, Eschbach PJ, et al. Are organic foods safer or healthier than conventional alternatives?: a systematic review. *Ann Intern Med* [Internet]. 2012 Oct 16 [cited 2022 May 2];157(5):348–66. Available from: <https://pubmed.ncbi.nlm.nih.gov/22944875/>
41. Crawley H, Westland S. [PDF] Infant milks in the UK: A practical guide for health professionals | Semantic Scholar [Internet]. 2012 [cited 2022 May 2]. Available from: <https://www.semanticscholar.org/paper/Infant-milks-in-the-UK-%3A-A-practical-guide-for-Trust/8da23275c1eb698667755912bd884e062f64d075>
42. Crawley H, Westland S. Fortified milks for children A worldwide review of fortified milks marketed for children over 1 year of age. 2013 [cited 2022 May 2]; Available from: www.firststepsnutrition.org
43. World Health Organization. First meeting of the WHO Scientific and Technical Advisory Group on Inappropriate Promotion of Foods for Infants and Young Children WHO headquarters, Geneva, Switzerland. 2013 [cited 2022 May 2]; Available from: <http://scalingupnutrition.org/>.
44. Baker P, Smith J, Salmon L, Friel S, Kent G, Iellamo A, et al. Global trends and patterns of commercial milk-based formula sales: is an unprecedented infant and young child feeding transition underway? *Public Health Nutr* [Internet]. 2016 Oct 1 [cited 2022 May 2];19(14):2540–50. Available from: <https://pubmed.ncbi.nlm.nih.gov/27211798/>